

Supplementary Table 6. Standardized mean differences with 95% confidence intervals from the network meta-analysis of intraoperative bleeding

Placebo	-0.256 (-0.587 to 0.076)	-0.577 (-0.829 to -0.325)	-1.055 (-1.161 to -0.881)	NA	NA
-0.437 (-0.650 to -0.225)/P=0.068	Beta-blocker	NA	0.423 (0.110 to 0.736)	NA	NA
-0.571 (-0.768 to -0.374)/P=0.405	-0.133 (-0.401 to 0.135)/NA	Clonidine	0.620 (0.140 to 1.100)	0.070 (-0.518 to 0.658)	NA
-0.971 (-1.161 to -0.781)/P=0.369	-0.534 (-0.741 to -0.327)/P=0.068	0.400 (0.173 to 0.628)/P=0.240	Dexmedetomidine	-0.238 (-0.599 to 0.123)	-0.026 (-0.533 to 0.480)
-0.691 (-0.968 to -0.414)/NA	-0.254 (-0.554 to 0.046)/NA	-0.121 (-0.402 to 0.161)/P=0.779	0.280 (0.051 to 0.503)/P=0.779	Opioid	NA
-0.945 (-1.420 to -0.470)/NA	-0.507 (-0.989 to -0.025)/NA	-0.374 (-0.865 to 0.117)/P=0.0684	0.026 (-0.409 to 0.462)/NA	-0.253 (-0.745 to 0.239)/NA	Nitroglycerine

Treatments are reported in alphabetical order. The results of direct comparisons are shown in the upper triangle, and the network meta-analysis results and local inconsistencies are shown in the lower triangle. For both, the estimates were obtained based on comparisons of the row-indicated treatments with the column-indicated treatments. NA, not available.